

Table 3. Testing Results in New York State Medicaid**Testing results for Teratogen Measures using New York State 2012 Medicaid Linked Mother Baby Records**

Measures	Deliveries	Number	Percent
A. Class X medications within the 9 months prior to delivery. (Lower is more safe)	102,399	323	0.32%
B. Class X Medications within the 12 months prior to delivery. (Lower is more safe)	102,399	1167	1.14%
C. Among High Risk pregnancies (pregnancy complications / maternal comorbidities) Class X Medications within 9 months prior delivery. (Lower is more safe)	61,713	323	0.43%
D. Among High Risk pregnancies (pregnancy complications/ maternal comorbidities plus teratogen use within 12 months of delivery) Class X Medications within 9 months prior delivery (Lower is more safe)	61,976	323	0.52%
E. Among High Risk pregnancies (pregnancy complications / maternal comorbidities) Class X Medications within 12 months prior to delivery. (Lower is more safe)	61,713	904	1.46%
F. Among High Risk pregnancies (pregnancy complications/ maternal comorbidities plus teratogen use within 12 months of delivery) Class X Medications within 12 months prior delivery. (Lower is more safe)	61,976	1167	1.88%
G. Women who have at least one refill of a teratogenic medication in the 15 months prior to pregnancy who have no prescriptions filled for teratogenic medication in the 9 months prior to delivery. (Higher is more safe)	984	803	81.61%